



The Commonwealth of Massachusetts
Executive Office of Health and Human Services
Department of Public Health
Bureau of Environmental Health
250 Washington Street, Boston, MA 02108-4619
Phone: 617-624-5757 Fax: 617-624-5777
TTY: 617-624-5286

CHARLES D. BAKER
Governor

KARYN E. POLITO
Lieutenant Governor

MARYLOU SUDDERS
Secretary

MONICA BHAREL, MD, MPH
Commissioner

Tel: 617-624-6000
www.mass.gov/dph

April 9, 2015

Susan G. Rask, M.S., R.S.
Public Health Director
Town of Concord
141 Keyes Road
Concord, MA 01742

Dear Ms. Rask:

Thank you for your letter of March 27, 2015, in which you requested that the Massachusetts Department of Public Health, Bureau of Environmental Health (MDPH/BEH), provide you with information concerning the use of crumb rubber infill material for artificial turf fields and potential health impacts. As you are likely aware, our office has had similar requests from some other municipalities regarding this issue, most recently from the health agent in Medway, Massachusetts.

In response to your request, I am including the summary provided to Medway on March 23, 2015, of our review of the most current scientific literature related to potential exposure to artificial turf components, including crumb rubber infill, and health impacts, including cancer, in relation to exposure to such turf. Recent local media reports on soccer players, particularly goalies, that have played on artificial turf and the incidence of some cancers have been expressed, notably non-Hodgkin's lymphoma, Hodgkin Lymphoma, and osteosarcoma.

Updated Literature Review

In previous evaluations performed during the period 2008-2013 for the Town of Needham, we noted that crumb rubber infill has been found to contain chemicals, including polycyclic aromatic hydrocarbons (PAHs), volatile organic compounds (VOCs), and metals. We further stated that although these chemicals are in the material itself, information available at that time did not suggest significant exposure opportunities to the chemicals in the materials such that we would expect health effects.

We noted that the most relevant study on this topic at the time was a study conducted by the California Office of Environmental Health Hazard Assessment (CA OEHHA).

Since that time, the CA OEHHA conducted additional evaluations of chemical concentrations in air above crumb rubber turf fields under active use (CA OEHHA 2010). Air samples were taken above fields and analyzed for VOCs and metals. Results suggested that adverse health effects were unlikely to occur from inhalation of VOCs or metals in particulates above these fields. To assess the potential for skin infections due to bacteria or to skin abrasions on these fields, tests for bacterial contamination were performed and the frequency of skin abrasions was assessed. Researchers found fewer bacteria detected on the artificial turf compared to natural turf, suggesting that the risk of infection to athletes using these fields was actually lower. However, more skin abrasions were observed in athletes using artificial turf fields than natural turf fields, and the study authors made various recommendations to help prevent skin abrasions (e.g., protective equipment or clothing) and prompt treatment of skin abrasions.

In another study, the state of Connecticut conducted air sampling at four outdoor artificial turf fields with crumb rubber infills (most relevant to Medway) under summer conditions (Simcox et al. 2011; Ginsberg et al. 2011). Air measurements were taken using stationary air sampling monitoring devices as well as personal samplers (placed on people using the fields). They concluded that exposure opportunities to turf contaminants were not associated with elevated health risks and suggested that their findings were consistent with other studies available at the time. A letter prepared by the Connecticut Department of Public Health reiterates these conclusions (CTDPH 2015).

A 2014 study by researchers at the Rutgers Robert Wood Johnson Medical School in New Jersey evaluated opportunities for exposures to PAHs, semivolatile organic compounds (SVOCs), and heavy metals from exposures to artificial turf fibers and crumb rubber infills by measuring these constituents in simulated body fluids (digestive fluids, lung fluids, sweat) that represented different routes of exposure (ingestion, inhalation, dermal). This bioaccessibility study aimed to provide a better measure of the actual amount of these contaminants that might be absorbed into the body after exposure. The researchers found that PAHs were routinely below the limit of detection and SVOCs that have environmental regulatory limits to use for comparison were identified at levels too low to quantify. Some metals were detected but at concentrations at which health risks were low, with the exception of lead from the field sample collected. That sample indicated lead at levels in the simulated digestive fluids that the authors reported could result in blood lead levels above the current U.S. Centers for Disease Control and Prevention (CDC) reference value for blood lead in children (5 ug/dL). It should be noted that the lead concentration of the materials used in this study included a sample of turf fiber with a lead concentration of 4,400 mg/kg. Based on the lead result from this one field sample, the authors suggested that components of artificial turf fields should be certified for low or no lead content prior to use. Overall, however, the authors concluded that opportunities for exposure to constituents in these fluids presented very low risk among all populations that would use artificial turf fields

(Pavilonis et al. 2014). Based on information you provided, it is our understanding that Concord will require artificial turf that will be installed to meet the Consumer Product Safety Improvement Act (CPSIA) limit of 100 mg/kg for lead in children's products and thus would be consistent with these recommendations.

A study conducted in 2010 in the Netherlands assessed the exposure of soccer players to PAHs after playing sports on a rubber crumb field. Urine testing in participants indicated that uptake of PAHs by the participants following exposure to artificial turf with rubber crumb infill was minimal. If there is any exposure, the authors reported, uptake is minimal and within the normal range of uptake of PAHs from environmental sources and/or diet observed in healthy individuals (van Rooij and Jongeneelen 2010).

It is probably worthwhile to also note that MDPH/BEH reviewed testing data for artificial turf for the Town of Needham, as reported in our letters of 2011 and 2013 to the Needham Board of Health. The Town of Needham contracted with an environmental testing firm to conduct environmental tests including, air measurements of volatile organic compounds taken in the laboratory and heavy metals (arsenic, cadmium, chromium, lead, mercury, selenium, zinc) content of crumb rubber materials. Our review and conclusions for that testing, did not indicate exposures of health concern.

Concerns About Cancer Among Soccer Players

As noted earlier in this letter, some recent news reports suggested that the incidence of cancers among soccer players, particularly goaltenders exposed to artificial turf, might be atypical. These media reports included many cancer types, but some focused specifically on NHL, Hodgkin Lymphoma, and osteosarcoma. We thought it would be helpful to provide additional information on cancers in general and known risk factors for NHL, Hodgkin Lymphoma, and osteosarcoma.

Cancer in General

Understanding that cancer is not one disease, but a group of diseases, is very important. Research has shown that there are more than 100 different types of cancer, each with separate causes, risk factors, characteristics and patterns of survival. A risk factor is anything that increases a person's chance of developing cancer and can include hereditary conditions, medical conditions or treatments, infections, lifestyle factors, or environmental exposures. Although risk factors can influence the development of cancer, most do not directly cause cancer. An individual's risk for developing cancer may change over time due to many factors and it is likely that multiple risk factors influence the development of most cancers. In addition, an individual's risk may depend on a complex interaction between their genetic make-up and exposure to environmental agents, including infectious agents and/or chemicals. This may explain why some individuals have a fairly low risk of developing a particular type of cancer as a result of an environmental exposure, while others are more vulnerable.

Cancers in general have long latency or development periods that can range from 10 to 30 years in adults, particularly for solid tumors. In some cases, the latency period may be more than 40 to 50 years. It is important to note, however, that latency periods for children and adolescents are significantly shorter than for adults.

Hodgkin Lymphoma

Hodgkin Lymphoma is most common in young adults between the ages of 15 and 40, especially in individuals in their 20s. Among adolescents, it is the most common type of cancer.

Hodgkin Lymphoma occurs specifically in a type of B lymphocyte (or white blood cell) called the Reed-Sternberg cell while other lymphomas (non-Hodgkin's types) occur in different cells.

Established risk factors for Hodgkin Lymphoma include: exposure to the Epstein-Barr virus (EBV); a previous diagnosis of mononucleosis (mono is caused by the EBV); family history; and certain hereditary conditions (such as ataxia telangiectasia) associated with a weakened immune system. The Epstein-Barr virus is very prevalent in the general population. Even though most of us have been exposed to the virus (which remains latent in our bodies), most people do not develop mononucleosis or Hodgkin Lymphoma. EBV is thought to account for about 20% or 25% of the diagnoses of classical Hodgkin's in the US.

Higher socioeconomic status is also a possible risk factor. This is thought to be due to delayed infectious exposures in childhood.

Occupational exposures as risk factors have been studied extensively and none have emerged as established risk factors. Likewise, there is very little evidence linking the risk of Hodgkin Lymphoma to an environmental exposure, other than the EBV.

Non-Hodgkin Lymphoma (NHL)

NHL refers to a diverse group of cancers that are characterized by an increase in malignant cells of the immune system. Each subtype of NHL may have different risk factors associated with its development. The specific cause of NHL in most individuals is unknown.

Although some types of NHL are among the more common childhood cancers, more than 95% of diagnoses occur in adults. Incidence generally increases with age, and most diagnoses occur in people in their 60s or older.

Established risk factors for NHL include a weakened immune system, associated with various medical conditions, and exposure to various viruses. An increased risk is faced by individuals taking immunosuppressant drugs following organ transplants; individuals with autoimmune disorders, such as rheumatoid arthritis and lupus; and individuals who

have taken certain chemotherapy drugs for other cancers. Several viruses have been shown to play a role in the development of NHL, including the human immunodeficiency virus (HIV), the human T-cell leukemia/lymphoma virus (HTLV-1), and the Epstein-Barr virus.

Exposure to high-dose radiation (for example, by survivors of atomic bombs and nuclear reactor accidents and possibly by patients who have received radiation therapy for a previous cancer) may pose an increased risk. Some studies have also suggested that exposure to chemicals such as benzene and certain herbicides and insecticides may be linked with an increased risk of NHL. Smoking has been associated in some studies with certain types of NHL.

Osteosarcoma

Osteosarcoma is a type of malignant bone cancer which accounts for about 2% of childhood cancers in the United States. It is the most common type of cancer that develops in bone and comprises about 66% of malignant bone tumors in children in Massachusetts. Most osteosarcomas occur in children and young adults between the ages of 10 and 30. Teenagers comprise the most commonly affected age group and are at the highest risk during their growth spurt. However, osteosarcoma can occur in people of any age, with about 10% of all osteosarcomas occurring in people over the age of 60.

Established risk factors for osteosarcoma include certain inherited syndromes (such as retinoblastoma, the Li-Fraumeni syndrome, and others) and certain bone diseases (such as Paget disease of the bone and hereditary multiple osteochondromas). Individuals with these syndromes and bone diseases have an increased risk of developing osteosarcoma. People who have received radiation treatment for a previous cancer may have a higher risk of later developing osteosarcoma in the area that was treated. Being treated at a younger age and with higher doses of radiation both increase the risk. Because the risk of osteosarcoma is highest between the ages of 10 and 30, especially during the teenage growth spurt, experts believe that there may be a link between rapid bone growth and the risk of a bone tumor. Children with osteosarcoma are often tall for their age, which supports the link with rapid bone growth. Other than radiation, there are no known lifestyle or environmental risk factors associated with osteosarcoma. Aside from these risk factors, the causes of most osteosarcomas are unknown.

In summary, the scientific literature continues to suggest that exposure opportunities to artificial turf fields are not generally expected to result in health effects. Use of artificial turf materials that meet CPSIA limits for lead content in children's products would minimize exposure opportunities to lead.


With respect to your report that Concord will require some testing of the artificial turf materials in the future, we would be happy to assist you and the Town of Concord in

developing a sampling and analysis plan as well as provide technical support in interpreting results, similar to the assistance that we provided to the Town of Needham.

As we stated in previous letters to municipalities on artificial turf, while available information does not indicate exposure opportunities of health concern, MDPH/BEH continues to recommend common sense ways to minimize any potential exposure to chemicals that may be contained in synthetic turf fields made of crumb rubber. MDPH/BEH suggests washing hands after playing on the field and before eating, particularly for younger children with frequent hand-to-mouth activity, and taking off shoes before entering the house to prevent tracking in any crumb rubber particles. Also, there are studies that indicate heat levels on artificial turf fields may rise as outdoor temperatures increase (New York State 2009). Thus, for protection of the players, MDPH/BEH recommends increasing hydration, taking frequent breaks, and watering down the field to cool it on hot days to prevent the potential for burns or heat stress. Finally, based on recent work in California, MDPH/BEH recommends that steps be taken to minimize the potential for skin abrasions (e.g., protective equipment) and that skin abrasions be treated promptly to prevent potential infections.

We hope this information is helpful to you and Concord residents. If you have any questions, please feel free to contact us at 617-624-5757.

Sincerely,



Suzanne K. Condon, Associate Commissioner
Director, Bureau of Environmental Health

References

American Cancer Society. 2015a. Detailed Guide: Hodgkin disease. Available at <http://www.cancer.org/cancer/hodgkindisease/detailedguide/index>. Last updated March 4.

American Cancer Society. 2015b. Detailed Guide: Non-Hodgkin lymphoma. Available at <http://www.cancer.org/cancer/non-hodgkinlymphoma/detailedguide/index>. Last updated March 11.

American Cancer Society. 2015c. Detailed Guide: Osteosarcoma. Available at <http://www.cancer.org/cancer/osteosarcoma/detailedguide/index>. Last updated January 6.

California Office of Environmental Health Hazard Assessment. 2010. Safety Study of Artificial Turf Containing Crumb Rubber Infill Made from Recycled Tires: Measurements of Chemicals and Particulates in the Air, Bacteria in the Turf, and Skin Abrasions Caused by Contact with the Surface. OEHHA, Pesticide and Environmental Toxicology

Branch, Funded by the Department of Resources Recycling and Recovery. October 2010, 121 p.

Connecticut Department of Public Health. 2015. Recent News Concerning Artificial Turf Fields. Letter to Local Health Departments and Districts, January 20, 2015. Connecticut Department of Public Health, Hartford, CT.

Gisberg, G; B. Toal; N. Simcox; A. Bracker; B. Bolembiewski; T. Kurland; C. Hedman. 2011. Human Health Risk Assessment of Synthetic Turf Fields Based Upon Investigation of Five Fields in Connecticut. *J. Tox Environ. Health, Part A, Current Issues*. 74:17, 1150-1174.

New York State Department of Environmental Conservation and New York State Department of Health. 2009. An Assessment of Chemical Leaching, Releases to Air and Temperature at Crumb-Infilled Synthetic Turf Fields.

Pavilonis, BT; CP Weisel; B. buckley; and PJ Liroy. 2014. Bioaccessibility and Risk of Exposure to Metals and SVOCs in Artificial Turf Field Fill Materials and Fibers. *Risk Anal*. 34:44-55.

Simcox, NJ; A Bracker; G. Ginsberg; B Toal; B. Golembiewski; T. Kurland; and C. Hedman. 2011. Synthetic Turf Field Investigation in Connecticut. *J Tox Environ Health, Part A*: 74(17):1133-1149.

Synthetic Turf Council. 2014. Guidelines for Crumb Rubber Infill Used in Synthetic Turf Fields. Printed October 2010, Revised October 23, 2014. Atlanta, GA.

Ulirsch, G; K Gleason; S. Gerstenberger; D Moffett; G. Pulliam, T ahmed; and J. Fagliano. 2010. Evaluating and Regulating Lead in Synthetic Turf. *Environ. Health Perspect.*, 118:1345-1349.

von Rooij, DJ, and PJ Jorgeneelen. 2010. Hydroxypyrene in urine of football players after playing on artificial sports field with the crumb rubber infill. *Int Arch Occup Environ Health*, 83(1):105-110. DOI: 10.1007/s00420-009-0465-y.